

# Determinants

## One Shot

$$D = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$



# Nishant Vora

**B.Tech - IIT Patna**

- 7+ years Teaching experience
- Mentored 5 lac+ students
- Teaching Excellence Award

# B<sup>o</sup>unceBa<sup>o</sup>ck




# Join with us in Telegram

## TELEGRAM CHANNEL

- [t.me/unacademyatoms](https://t.me/unacademyatoms)

## COMPLETE NOTES AND LECTURES

- [tinyurl.com/unacademyatoms](https://tinyurl.com/unacademyatoms)



# + India's **BEST** Educators

# Unacademy Subscription

**BATCH**

**HINDI**

Igneous Batch for JEE Advanced & Olympiads 2021

Started on Jul 7

Nishant Vora and 1 more

**BATCH**

**HINDI**

Evolve Batch Course for Class 12th JEE Main and Advanced 2022

Starts on Apr 7

Anupam Gupta and 2 more

**BATCH**

**HINDI**

Mega Batch Course for Class 12th JEE Main and Advanced 2022

Starts on Apr 6

Narendra Avasthi and 1 more

**BATCH**

**HINDI**

Enthuse: Class 12th for JEE Main and Advanced 2022

Starts on Apr 14

Amarnath Anand and 2 more

**BATCH**

**HINDI**

Final Rapid Revision Batch for JEE Main 2021

Starts on Apr 6

Manoj Chauhan and 2 more

**plus**






**HINDI PHYSICS**

Course of 12th syllabus Physics for JEE Aspirants 2022: Part - I

Lesson 1 • Apr 2, 2021 12:30 PM

D C Pandey

If you want to be the **BEST**  
“Learn” from the **BEST**





×

IIT JEE subscription

ICONIC <sup>+</sup> PLUS

✓ India's Best Educators

✓ Interactive Live Classes

✓ Structured Courses & PDFs

✓ Live Tests & Quizzes

✗ 1:1 Live Mentorship

✗ Live Doubt Solving

30 months

₹1,980/mo

No cost EMI

+10% OFF ₹59,399

>

24 months

₹2,310/mo

No cost EMI

+10% OFF ₹55,440

>

12 months

₹3,176/mo

No cost EMI

+10% OFF ₹38,115

>

6 months

₹4,620/mo

No cost EMI

+10% OFF ₹27,720

>

3 months

₹5,775/mo

+10% OFF ₹17,325

>

To be paid as a one-time payment

NVLIVE

×



×

IIT JEE subscription

PLUS ICONIC <sup>+</sup>

✓ India's Best Educators

✓ Interactive Live Classes

✓ Structured Courses & PDFs

✓ Live Tests & Quizzes

✓ Personal Coach

✓ Study Planner

24 months

₹3,750/mo

No cost EMI

+10% OFF ₹90,000

>

18 months

₹4,000/mo

No cost EMI

+10% OFF ₹72,000

>

12 months

₹4,875/mo

No cost EMI

+10% OFF ₹58,500

>

6 months

₹5,700/mo

No cost EMI

+10% OFF ₹34,200

>

To be paid as a one-time payment

NVLIVE

×






# Determinant



# Determinants





# Representation



$$\underline{|A|} = \begin{vmatrix} \mathbf{a} & \mathbf{b} \\ \mathbf{c} & \mathbf{d} \end{vmatrix}_{2 \times 2} = \underline{\det(A)}$$

$$\underline{|A| = \det(A)}$$

1x1
2x2
3x3
4x4





## Determinant value of $1 \times 1$ & $2 \times 2$

$$A = \begin{vmatrix} -2 \end{vmatrix}_{1 \times 1} = (-2)$$

$$A = \begin{vmatrix} a & b \\ c & d \end{vmatrix}_{2 \times 2} = \boxed{ad - bc}$$



The value of  $\left| \begin{array}{cc} a+1 & a-2 \\ a+2 & a-1 \end{array} \right|$  is

A.  $2a^2$

B. 0

C. -3

☒ D. 3

$$(a+1)(a-1) - (a-2)(a+2)$$

$$= (\cancel{a^2} - 1) - (\cancel{a^2} - 4)$$

$$= \textcircled{3}$$



The value of  $\left| \frac{1 + \cos\theta}{\sin\theta} \times \frac{\sin\theta}{1 - \cos\theta} \right|$  is

A. 2

B. -1

C. 0

D.  $\cos 2\theta$

$$\Rightarrow (1 + \cos\theta)(1 - \cos\theta) - \sin^2\theta$$

$$\Rightarrow (1 - \cos^2\theta) - \sin^2\theta$$

$$\Rightarrow 0$$



# Minor & Cofactor



# Minors

$$D = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}_{3 \times 3}$$

$$M_{11} = \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix}$$

$$M_{21} = \begin{vmatrix} a_{12} & a_{13} \\ a_{32} & a_{33} \end{vmatrix}$$

$$M_{32} = \begin{vmatrix} a_{11} & a_{13} \\ a_{21} & a_{23} \end{vmatrix}$$





## Cofactor :

$$C_{ij} = (-1)^{i+j} M_{ij}$$

$$D = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

$$C_{11} = (-1)^{1+1} M_{11} = M_{11}$$

$$C_{12} = (-1)^{1+2} M_{12} = -M_{12}$$

$$C_{13} = (-1)^{1+3} M_{13} = M_{13}$$

$$C_{ij} = (-1)^{i+j} M_{ij}$$

$$\underline{i+j = \text{odd}}$$

$$C_{12} = -M_{12}$$

$$C_{21} = -M_{21}$$

$$C_{32} = -M_{32}$$

$$C_{23} = -M_{23}$$



**Remember :**

$$C_{ij} = (-1)^{i+j} M_{ij}$$

$$D = \begin{vmatrix} \mathbf{a}_{11} & \mathbf{a}_{12} & \mathbf{a}_{13} \\ \mathbf{a}_{21} & \mathbf{a}_{22} & \mathbf{a}_{23} \\ \mathbf{a}_{31} & \mathbf{a}_{32} & \mathbf{a}_{33} \end{vmatrix}$$

$$D = \begin{vmatrix} + & \boxed{-} & + \\ \boxed{-} & + & \boxed{-} \\ + & \boxed{-} & + \end{vmatrix}$$





# Expanding/Opening Determinant

Expanding w.r.t R1

$$D = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

w.r.t R1

$$\Rightarrow a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13}$$

w.r.t  $C_2$

$$\Rightarrow a_{12}C_{12} + a_{22}C_{22} + a_{32}C_{32}$$





## Determinant value of $3 \times 3$

$$A = \begin{vmatrix} 2 & -3 & 1 \\ 2 & 0 & -1 \\ 1 & 4 & 5 \end{vmatrix}$$

Minor

$$\begin{vmatrix} 4 & 11 & 8 \\ -19 & 9 & 11 \\ 3 & -4 & 6 \end{vmatrix}$$

Cofactor

$$\begin{vmatrix} 4 & -11 & 8 \\ 19 & 9 & -11 \\ 3 & 4 & 6 \end{vmatrix}$$

w.r.t R1

$$(2)(4) + (-3)(-11) + (1)(8) = \boxed{49}$$

w.r.t C<sub>2</sub>

$$(-3)(-11) + (0)9 + (4)4 = \boxed{49}$$



## Determinant value of 3×3



$$\begin{vmatrix} 2 & -3 & 1 \\ 2 & 0 & -1 \\ 1 & 4 & 5 \end{vmatrix}$$

$$(2)(3) + (-3)(4) + (1)(6) \Rightarrow \boxed{0}$$

$$(-3)(8) + (0)(-11) + (4)(6) = \boxed{0}$$



## Cofactor property

In a determinant **the sum of the product's** of the element's of any row (column) with their corresponding cofactor's is **equal to the value of determinant**.

$$D = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$



## Cofactor property



$$\begin{vmatrix} 2 & -3 & 1 \\ 2 & 0 & -1 \\ 1 & 4 & 5 \end{vmatrix}$$



**Shortcut  
to find value of  
determinant**



## #Shortcut (Rule of Sarrus)



$$\begin{vmatrix} a & b & c \\ e & f & g \\ h & i & j \end{vmatrix} = \begin{vmatrix} + & + & + \\ a & b & c \\ e & f & g \\ h & i & j \end{vmatrix} \begin{vmatrix} a & b \\ e & f \\ h & i \end{vmatrix} = afj + bgh + cei - hfc - iga - jeb$$

The diagram illustrates the Rule of Sarrus for calculating the determinant of a 3x3 matrix. It shows the original matrix, the expanded matrix with signs, and the final result. Blue solid arrows indicate the positive terms (afj, bgh, cei) and red dashed arrows indicate the negative terms (hfc, iga, jeb).



## #Shortcut (Rule of Sarrus)



2	-3	1
2	0	-1
1	4	5

Diagram illustrating the Rule of Sarrus for a 3x3 matrix. The first two columns are repeated to the right. Purple arrows show the downward diagonal products (2\*0\*5, -3\*5\*1, 1\*1\*2) and upward diagonal products (1\*0\*1, -1\*1\*2, 5\*2\*-3). Red numbers indicate the products: 0, -3, 5, 1, -2, -15.

$$(0 + 3 + 8) - (0 - 8 - 30) = \boxed{49}$$

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix}$$

$$ad - bc$$



## #Shortcut (Rule of Sarrus)

$$\begin{bmatrix} 2 & 3 & 1 \\ 1 & 1 & 1 \\ 2 & 2 & 1 \end{bmatrix}$$

Diagram illustrating the Rule of Sarrus for a 3x3 matrix. The matrix is shown with its elements. Blue arrows indicate the forward diagonal products (down-right):  $2 \cdot 1 \cdot 1 = 2$ ,  $3 \cdot 1 \cdot 1 = 3$ , and  $1 \cdot 2 \cdot 2 = 4$ . Green arrows indicate the backward diagonal products (down-left):  $1 \cdot 1 \cdot 2 = 2$ ,  $1 \cdot 1 \cdot 1 = 1$ , and  $2 \cdot 2 \cdot 3 = 12$ . The final result is calculated as  $(2 + 3 + 4) - (2 + 1 + 12) = 9 - 15 = -6$ .

$$(2 + 6 + 2) - (2 + 4 + 3) \\ = \boxed{1}$$





# Properties of Determinant



# Properties of Determinants



$$\textcircled{1} \quad |A^T| = |A|$$

$$\textcircled{2} \quad |I| = 1$$

$$1. \quad |A^T| = |A|$$

$$\text{Note: } |I| = 1$$

$$\text{Ex } \begin{vmatrix} 2 & 3 \\ 4 & 5 \end{vmatrix} = \begin{vmatrix} 2 & 4 \\ 3 & 5 \end{vmatrix}$$
$$10 - 12 = 10 - 12$$



# Properties of Determinants

2. If any two rows (or columns) of a determinant **be interchanged**, the value of determinant is **changed in sign only**.

$$\begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix} = - \begin{vmatrix} p & q & r \\ a & b & c \\ x & y & z \end{vmatrix}$$



## Properties of Determinants

3. If row or columns are rotated in cyclic order then value of determinant is **unchanged**

→ Same

$$\begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix} = \begin{vmatrix} x & y & z \\ a & b & c \\ p & q & r \end{vmatrix}$$

1 Cycle = 2 Swaps

$$- \begin{vmatrix} p & q & r \\ a & b & c \\ x & y & z \end{vmatrix}$$



# Properties of Determinants



4. If a determinant has **any two rows (or columns) identical**, then its **value is zero**.

$$\begin{vmatrix} a & b & c \\ a & b & c \\ x & y & z \end{vmatrix} = 0$$

$$x(0) + y(0) + z(0) = 0$$

1 2

$$\begin{vmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ x & y & z \end{vmatrix} = 0$$

$$2 \begin{vmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \\ x & y & z \end{vmatrix} = 0$$



# Properties of Determinants



**5. Scalar multiplication:** Scalar will be multiplied in any one row (or column)

**e.g.** If  $D = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$  then  $kD$  =  $\begin{vmatrix} ka_1 & kb_1 & kc_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$

$$k \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} ka & kb \\ kc & kd \end{bmatrix}$$



# Properties of Determinants



★★

6.  $|kA| = k^n |A|$ , where  $n$  is order of  $A$ .

$$|kA| = k^n |A|$$

Diagram illustrating the property: A red arrow points from the 'k' in  $|kA|$  to the 'n' in  $k^n$ . Two purple arrows point from the 'A' in  $|kA|$  to the 'A' in  $|A|$ .

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$kA = \begin{bmatrix} ka & kb \\ kc & kd \end{bmatrix}$$

$$\begin{aligned} |kA| &= \begin{vmatrix} ka & kb \\ kc & kd \end{vmatrix} \\ &= k^2 \begin{vmatrix} a & b \\ c & d \end{vmatrix} \end{aligned}$$



Evaluate

102	18	36
1	3	36
17	3	6

= 6

$$\begin{vmatrix} 17 & 3 & 6 \\ 1 & 3 & 36 \\ 17 & 3 & 6 \end{vmatrix} = 0$$

$$\underline{6 \quad 1}$$







Let  $A = [a_{ij}]$  and  $B = [b_{ij}]$  be two  $3 \times 3$  real matrices such that

$b_{ij} = (3)^{(i+j-2)} a_{ij}$ , where  $i, j = 1, 2, 3$ . If the determinant of B is 81, then

the determinant of A is :

A.  $1/3$

B.  $3$

C.  $1/81$

☒ D.  $1/9$

$$b_{ij} = 3^{i+j-2} a_{ij}$$

(Given)


$$|B| = 81$$

*JEE Main 2020*

$$|A| = ?$$

$$B = \begin{bmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \\ b_{31} & b_{32} & b_{33} \end{bmatrix}_{3 \times 3}$$

$$|B| = \begin{vmatrix} a_{11} & 3a_{12} & 3^2a_{13} \\ 3a_{21} & 3^2a_{22} & 3^3a_{23} \\ 3^2a_{31} & 3^3a_{32} & 3^4a_{33} \end{vmatrix} = 81$$



$$3^3 \begin{vmatrix} a_{11} & \boxed{\cancel{3} a_{12}} & \boxed{\cancel{3^2} a_{13}} \\ a_{21} & \boxed{\cancel{3} a_{22}} & \boxed{\cancel{3^2} a_{23}} \\ a_{31} & \boxed{\cancel{3} a_{32}} & \boxed{\cancel{3^2} a_{33}} \end{vmatrix} = 81$$

$$3^3 \cdot 3^3 |A| = 81$$

$$|A| = \frac{1}{9}$$



Let  $A$  be a  $3 \times 3$  matrix with  $\det(A) = 4$ . Let  $R_i$  denote the  $i^{\text{th}}$  row of  $A$ . If a matrix  $B$  is obtained by performing the operation  $R_2 \rightarrow 2R_2 + 5R_3$  on  $2A$ , then  $\det(B)$  is equal to

*JEE Main 2021*

- ☒ A. 64
- ☐ B. 16
- ☐ C. 80
- ☐ D. 128

$$|A| = 4 \quad |B| = ?$$

$$2A = \begin{bmatrix} 2a & 2b & 2c \\ 2x & 2y & 2z \\ 2p & 2q & 2r \end{bmatrix} \quad 2(R_2) + 5R_3$$

$$|B| = \begin{vmatrix} 2a & 2b & 2c \\ 4x+10p & 4y+10q & 4z+10r \\ 2p & 2q & 2r \end{vmatrix}$$

$$|B| = \begin{vmatrix} 2a & 2b & 2c \\ 4x & 4y & 4z \\ 2p & 2q & 2r \end{vmatrix} + \begin{vmatrix} 2a & 2b & 2c \\ 10p & 10q & 10r \\ 2p & 2q & 2r \end{vmatrix} \rightarrow 0$$

$$= 16 \begin{vmatrix} a & b & c \\ x & y & z \\ p & q & r \end{vmatrix}$$

$$= 16 \times |A|$$

$$= 16 \times 4$$

$$= \textcircled{64}$$



Let p and p + 2 be prime numbers and let

$$\Delta = \begin{vmatrix} p! & (p+1)! & (p+2)! \\ (p+1)! & (p+2)! & (p+3)! \\ (p+2)! & (p+3)! & (p+4)! \end{vmatrix}$$



$\alpha_{\max} + \beta_{\max}$

Then the sum of the maximum values of  $\alpha$  and  $\beta$ , such that  $p^\alpha$  and  $(p+2)^\beta$  divide  $\Delta$ , is 4.


$$\Delta = p! (p+1)! (p+2)! \begin{vmatrix} 1 & p+1 & (p+2)(p+1) \\ 1 & p+2 & (p+3)(p+2) \\ 1 & p+3 & (p+4)(p+3) \end{vmatrix}$$

*JEE Main 2022*

$$R_2 \rightarrow R_2 - R_1$$

$$R_3 \rightarrow R_3 - R_2$$

$$\Delta = p! (p+1)! (p+2)! \begin{vmatrix} 1 & p+1 & (p+2)(p+1) \\ 0 & 1 & 2p+4 \\ 0 & 1 & 2p+6 \end{vmatrix}$$



$$p^{\alpha=3} (p+2)^{\beta=1}$$

$$\Delta = \underline{p!} \underline{(p+1)!} \underline{(p+2)!} \times 2$$

$$= \underline{p} (p-1)! (p+1)(p) (p-1)! \underline{(p+2)} (p+1) p (p-1)! \times 2$$



# Properties of Determinants



Note: The value of a **skew symmetric** determinant of **odd order** is zero.

$$\begin{vmatrix} 0 & a & -b \\ -a & 0 & c \\ b & -c & 0 \end{vmatrix}$$

The determinant is shown with red lines crossing out the diagonal elements (0, 0, 0) and connecting the off-diagonal elements (a, -b, c, -c, b, -a) in a cycle, illustrating the skew-symmetric property.

\*\*\*\*\*

Skew-sym. + odd order

$$|A| = 0$$

- ① all diagonal elements must be "0"
- ② Mirror Image  $\Rightarrow$  Sign will be opp



# Properties of Determinants



## 7. Adding Determinants

$$\begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix} + \begin{vmatrix} x & y & z \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix} = \begin{vmatrix} a_1 + x & b_1 + y & c_1 + z \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

$$\begin{vmatrix} - & - & 1 \\ - & - & 2 \\ - & - & 3 \end{vmatrix} + \begin{vmatrix} - & - & 4 \\ - & - & 5 \\ - & - & 6 \end{vmatrix}$$

One at a time





# Properties of Determinants



## 8. Splitting Determinants

$$\begin{vmatrix} \underline{a_1 + x} & \underline{b_1 + y} & \underline{c_1 + z} \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix} = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix} + \begin{vmatrix} x & y & z \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

One at a time



Find  $\begin{vmatrix} a & b & c \\ a+2x & b+2y & c+2z \\ x & y & z \end{vmatrix}$

$$= \begin{vmatrix} a & b & c \\ a & b & c \\ x & y & z \end{vmatrix} + 2 \begin{vmatrix} a & b & c \\ x & y & z \\ x & y & z \end{vmatrix}$$

$$= 0 + 0$$

$$= \boxed{0}$$





# Properties of Determinants

9.  $|AB| = |A| |B|$

$$4 = 1 \times 4$$

$$\underline{|AB| = |A| |B|}$$

$$A = \begin{vmatrix} 1 & 1 \\ 1 & 2 \end{vmatrix} \text{ and } B = \begin{vmatrix} 1 & 0 \\ 1 & 4 \end{vmatrix} \text{ Then } AB = \begin{vmatrix} 2 & 4 \\ 3 & 8 \end{vmatrix}$$

$$\underline{|A| = 1}$$

$$\underline{|B| = 4}$$

$$AB = \begin{vmatrix} \boxed{1} & \boxed{1} \\ \boxed{1} & \boxed{2} \end{vmatrix}$$

$$\begin{vmatrix} \boxed{1} & \boxed{0} \\ \boxed{1} & \boxed{4} \end{vmatrix}$$

$$\underline{|ABC| = |A| |B| |C|}$$

$$|AB| = \begin{vmatrix} 2 & 4 \\ 3 & 8 \end{vmatrix} = 16 - 12 = 4$$



# Properties of Determinants



10. If  $\det(A) = 0$ , then A is known as **singular** matrix.

eg.  $\begin{vmatrix} 1 & 1 \\ 2 & 2 \end{vmatrix} = 0$   $\begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$

Singular Matrix

11)  $|A^n| = |A|^n$



Let  $\beta$  be a real number. Consider the matrix

$$I + A = \begin{pmatrix} \beta & 0 & 1 \\ 2 & 1 & -2 \\ 3 & 1 & -2 \end{pmatrix} + \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

If  $A^7 - (\beta - 1)A^6 - \beta A^5$  is a singular matrix, then the value of  $9\beta$  is \_\_\_\_\_.

$$|A^7 - (\beta - 1)A^6 - \beta A^5| = 0$$

$$|A| = \beta(0) + 1(-1)$$

*JEE Adv. 2022*

$$|A^5 (A^2 - (\beta - 1)A - \beta I)| = 0$$

$$|A| = -1$$

$$|A^5 (\underline{A^2 - \beta A} + \underline{A - \beta I})| = 0$$

$$|A + I| = \begin{vmatrix} \beta+1 & 0 & 1 \\ 2 & 2 & -2 \\ 3 & 1 & -1 \end{vmatrix}$$

$$|A^5 (A + I)(A - \beta I)| = 0$$

$$\begin{aligned} &= (\beta+1)(0) + 1(2-6) \\ &= -4 \end{aligned}$$

$$\underbrace{|A|}^5 \underbrace{|A+I|}_{\text{non-zero}} \underbrace{|A-\beta I|}_{\text{non-zero}} = 0$$

non-zero non-zero

$$A - \beta I = \begin{bmatrix} \beta & 0 & 1 \\ 2 & 1 & -2 \\ 3 & 1 & -2 \end{bmatrix} - \begin{bmatrix} \beta & 0 & 0 \\ 0 & \beta & 0 \\ 0 & 0 & \beta \end{bmatrix}$$

$$|A - \beta I| = \begin{vmatrix} 0 & 0 & 1 \\ 2 & 1-\beta & -2 \\ 3 & 1 & -2-\beta \end{vmatrix} = 0$$

$$2 - 3(1 - \beta) = 0$$

$$\beta = \frac{1}{3}$$

$$\boxed{9\beta = 3} \text{ Ans}$$



# Elementary Transformation



11. The value of determinant remains same if we apply elementary transformation

$$R_1 \rightarrow R_1 + kR_2 + mR_3 \text{ or } C_1 \rightarrow C_1 + kC_2 + mC_3$$

Row transform

$$\underline{R_1 \rightarrow 1R_1 + aR_2 + bR_3}$$

$a, b \in \text{Constants}$

$$R_3 \rightarrow 1R_3 + \underbrace{2R_1 - 3R_2}$$



Prove that

$$\begin{vmatrix} a & a+b & a+b+c \\ 2a & 3a+2b & 4a+3b+2c \\ 3a & 6a+3b & 10a+6b+3c \end{vmatrix} = \underline{a^3}$$

Objective

①

○

○

~~□~~

②

1

1

1



$$3a + \cancel{2b} - 2(a + \cancel{b})$$

$$R_2 \rightarrow R_2 - 2R_1$$

$$R_3 \rightarrow \underline{R_3 - 3R_1}$$

$$\begin{vmatrix} a & a+b & a+b+c \\ 0 & a & 2a+b \\ 0 & 3a & 7a+3b \end{vmatrix}$$

$$\begin{aligned} &= a(7a^2 + \cancel{3ab} - 6a^2 - \cancel{3ab}) \\ &= \textcircled{a^3} \end{aligned}$$







The maximum value of  $f(x) = \begin{vmatrix} \frac{\sin^2 x}{1 + \sin^2 x} & \frac{1 + \cos^2 x}{\cos^2 x} & \cos 2x \\ \frac{\sin^2 x}{\cos^2 x} & \frac{\cos^2 x}{\sin 2x} & \cos 2x \end{vmatrix}, x \in R$  is

A.  $\sqrt{7}$

B.  $3/4$

✓ C.  $\sqrt{5}$

D.  $5$

$C_1 \rightarrow C_1 + C_2$

$f(x) = \begin{vmatrix} \frac{2}{1} & \frac{1 + \cos^2 x}{\cos^2 x} & \frac{\cos 2x}{\sin 2x} \\ \frac{2}{\cos^2 x} & \frac{\cos^2 x}{\sin 2x} & \cos 2x \end{vmatrix}$

$R_1 \rightarrow R_1 - R_2$


$f(x) = 1 \cos 2x - 2 \sin 2x$

$\sqrt{1^2 + 2^2} = \sqrt{5}$

$\begin{vmatrix} 0 & 1 & 0 \\ 2 & \cos^2 x & \cos 2x \\ 1 & \cos^2 x & \sin 2x \end{vmatrix} = -1 (2 \sin 2x - \cos 2x)$

16th Mar, 2021 (shift 2)




$$-\sqrt{a^2+b^2} \leq a \sin \theta + b \cos \theta \leq \sqrt{a^2+b^2}$$





The solutions of the equation

$$\begin{vmatrix} \frac{1 + \sin^2 x}{\cos^2 x} & \frac{\sin^2 x}{1 + \cos^2 x} & \frac{\sin^2 x}{\cos^2 x} \\ \frac{1}{4 \sin 2x} & \frac{1}{4 \sin 2x} & 1 + 4 \sin 2x \end{vmatrix} = 0$$

A.  $\pi/12, \pi/6$

B.  $\pi/6, 5\pi/6$

C.  $5\pi/12, 7\pi/12$

☒ D.  $7\pi/12, 11\pi/12$



$$R_1 \rightarrow R_1 + R_2$$

18th Mar, 2021 (shift 1)

$$\begin{vmatrix} \underline{2} & \underline{2} & 1 \\ \underline{\cos^2 x} & \underline{1 + \cos^2 x} & \cos^2 x \\ \underline{4 \sin 2x} & \underline{4 \sin 2x} & 1 + 4 \sin 2x \end{vmatrix} = 0$$

$$C_1 \rightarrow C_1 - C_2$$

$$\begin{vmatrix} 0 & 2 & 1 \\ 0 & 1 + \cos^2 x & \cos^2 x \\ 0 & 4 \sin 2x & 1 + 4 \sin 2x \end{vmatrix} = 0$$

$$1(2 + 8 \sin^2 x - 4 \sin 2x) = 0$$

$$2 + 4 \sin 2x = 0$$

$$\sin 2x = -\frac{1}{2}$$

$$2x = \frac{7\pi}{6}, \frac{11\pi}{6}$$

$$x = \frac{7\pi}{12}, \frac{11\pi}{12}$$

$$\underline{S-1} \quad \sin(\alpha) = \frac{1}{2}$$

$$\alpha = \frac{\pi}{6}$$

<u>S-2</u>	S	A
	$\pi$	$\frac{\pi}{6}$

<u>S-3</u>	$\pi - \alpha$	$\alpha$
	$\pi + \frac{\pi}{6}$	$2\pi - \frac{\pi}{6}$



$$\text{Let } f(x) = \begin{vmatrix} \sin^2 x & -2 + \cos^2 x & \cos 2x \\ 2 + \sin^2 x & \cos^2 x & \cos 2x \\ \sin^2 x & \cos^2 x & 1 + \cos 2x \end{vmatrix}, x \in [0, \pi]$$

Then the maximum value of  $f(x)$  is equal to

*JEE Main 2021*

**HOMEWORK**



The total number of distinct  $x \in \mathbb{R}$  for which

$$\begin{vmatrix} x & x^2 & 1+x^3 \\ 2x & 4x^2 & 1+8x^3 \\ 3x & 9x^2 & 1+27x^3 \end{vmatrix} = 10 \quad \text{is}$$

2 sol<sup>n</sup>

$$x^3 \begin{vmatrix} 1 & 1 & 1+x^3 \\ 2 & 4 & 1+8x^3 \\ 3 & 9 & 1+27x^3 \end{vmatrix} = 10$$

$$x^3 \left[ \begin{vmatrix} 1 & 1 & 1 \\ 2 & 4 & 1 \\ 3 & 9 & 1 \end{vmatrix} + x^3 \begin{vmatrix} 1 & 1 & 1 \\ 2 & 4 & 8 \\ 3 & 9 & 27 \end{vmatrix} \right] = 10$$
$$x^3 (2 + 12x^3) = 10$$

[JEE Adv  
2016]



$$2x^3 + 12x^6 = 10$$

$$x^3 = t$$

$$2t + 12t^2 = 10$$

$$6t^2 + t - 5 = 0$$

$$6t^2 + 6t - 5t - 5 = 0$$

$$(6t - 5)(t + 1) = 0$$

$$t = \frac{5}{6}, -1$$

$$x^3 = \frac{5}{6}, -1$$

$$x = \left(\frac{5}{6}\right)^{1/3}, -1$$





$$\boxed{x^3 = 1} \begin{cases} 1 \\ \omega \\ \omega^2 \end{cases}$$

$$\begin{aligned} \star \quad \omega^3 &= 1 \\ \star \quad 1 + \omega + \omega^2 &= 0 \end{aligned}$$

W.F.E

$$\boxed{x^2 + x + 1 = 0} \begin{cases} \omega \\ \omega^2 \end{cases}$$

# Advance Ques

(Determinant + Complex Numbers)

$$x^3 - 1 = 0$$

$$\underbrace{(x-1)}_{x=1} \underbrace{(x^2+x+1)}_{\substack{x=\omega \\ x=\omega^2}} = 0$$

$$x=1$$

$$x=\omega$$

$$x=\omega^2$$



Let  $\omega$  be the complex number  $\cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3}$ . Then the number of

distinct complex numbers  $z$  satisfying  $\begin{vmatrix} z+1 & \omega & \omega^2 \\ \omega & z+\omega^2 & 1 \\ \omega^2 & 1 & z+\omega \end{vmatrix} = 0$  is equal to

$\rightarrow$  (1)

$$\Rightarrow z^3 = 0$$

$$\Rightarrow \boxed{z=0} \text{ only Soln}$$

*JEE Adv 2010*



Let  $\alpha$  and  $\beta$  be the roots of the equation  $x^2 + x + 1 = 0$ . Then for

$y \neq 0$  in  $\mathbb{R}$ ,  $\begin{vmatrix} y+1 & \alpha & \beta \\ \alpha & y+\beta & 1 \\ \beta & 1 & y+\alpha \end{vmatrix}$  is equal to:

$$\begin{cases} \omega = \alpha \\ \omega^2 = \beta \end{cases}$$

A.  $y(y^2 - 1)$

B.  $y(y^2 - 3)$

☒ C.  $y^3$

D.  $y^3 - 1$

$$\begin{vmatrix} y+1 & \omega & \omega^2 \\ \omega & y+\omega^2 & 1 \\ \omega^2 & 1 & y+\omega \end{vmatrix}$$

$$R_1 \rightarrow R_1 + R_2 + R_3$$

$$\Rightarrow y \begin{vmatrix} 1 & 1 & 1 \\ \omega & y+\omega^2 & 1 \\ \omega^2 & 1 & y+\omega \end{vmatrix}$$

[JEE M 2019]



$$\Rightarrow y \left\{ 1 \left( (y+\omega)(y+\omega^2) - y \right) - 1(y\omega) + 1(-\omega^2 y) \right\}$$

$$\Rightarrow y \left\{ y^2 + \cancel{\omega^2 y} + \cancel{\omega y} - \cancel{y\omega} - \cancel{\omega^2 y} \right\}$$

$$\Rightarrow \textcircled{y^3}$$





Let  $\omega$  be the complex cube root of unity with  $\omega \neq 1$  and  $P = [P_{ij}]$  be a  $n \times n$  matrix with  $p_{ij} = \omega^{i+j}$ . Then  $P^2 \neq O$ , when  $n =$

A. 57

☒ B. 55

☒ C. 58

☒ D. 56

B C D

$$p_{ij} = \omega^{i+j} \quad P^2 \neq O \quad n = ?$$

for  $n=2$

$$P = \begin{bmatrix} p_{11} & p_{12} \\ p_{21} & p_{22} \end{bmatrix}_{2 \times 2} = \begin{bmatrix} \omega^2 & 1 \\ 1 & \omega \end{bmatrix}$$

[JEE Adv 2013]

$$P^2 = \begin{bmatrix} \omega^2 & 1 \\ 1 & \omega \end{bmatrix} \cdot \begin{bmatrix} \omega^2 & 1 \\ 1 & \omega \end{bmatrix} = \begin{bmatrix} \omega+1 & - \\ - & - \end{bmatrix} \neq O$$

for  $n=3$

$$p = \begin{bmatrix} p_{11} & p_{12} & p_{13} \\ p_{21} & p_{22} & p_{23} \\ p_{31} & p_{32} & p_{33} \end{bmatrix}_{3 \times 3} = \begin{bmatrix} \omega^2 & 1 & \omega \\ 1 & \omega & \omega^2 \\ \omega & \omega^2 & 1 \end{bmatrix}$$

$$p^2 = \begin{bmatrix} \omega^2 & 1 & \omega \\ 1 & \omega & \omega^2 \\ \omega & \omega^2 & 1 \end{bmatrix} \begin{bmatrix} \omega^2 & 1 & \omega \\ 1 & \omega & \omega^2 \\ \omega & \omega^2 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$n \rightarrow$  mult of 3  $p^2 = 0$

$n \rightarrow$  not " " "  $p^2 \neq 0$



Let  $\omega \neq 1$  be the cube root of unity and S be the set of all

non-singular matrices of the form 
$$\begin{bmatrix} 1 & a & b \\ \omega & 1 & c \\ \omega^2 & \omega & 1 \end{bmatrix}$$

where each of a, b and c is either  $\omega$  or  $\omega^2$ . Then the number of distinct matrices in the set S is

$a, b, c \in \{\omega, \omega^2\}$  [JEE Adv 2011]

A. 2

B. 6

C. 4

D. 8

$$\begin{vmatrix} 1 & a & b \\ \omega & 1 & c \\ \omega^2 & \omega & 1 \end{vmatrix} \neq 0$$

$$1(1-c\omega) - a(\omega - c\omega^2) + b(0) \neq 0$$

$$(1-c\omega) - a\omega(1-c\omega) \neq 0$$

$$(1-a\omega)(1-c\omega) \neq 0$$





$$(1 - aw)(1 - cw) \neq 0$$

↑                      ↑

$a \neq w^2$                $c \neq w^2$

	a	b	c
①	w	w	w
②	w	w <sup>2</sup>	w

Ans:- 2



If  $A = \begin{pmatrix} 0 & \sin \alpha \\ \sin \alpha & 0 \end{pmatrix}$  and  $\det(A^2 - \frac{1}{2}I) = 0$ , then a possible value of  $\alpha$  is

A.  $\pi/2$

B.  $\pi/3$

☒ C.  $\pi/4$

D.  $\pi/6$

$$A^2 = \begin{bmatrix} 0 & \sin \alpha \\ \sin \alpha & 0 \end{bmatrix} \begin{bmatrix} 0 & \sin \alpha \\ \sin \alpha & 0 \end{bmatrix}$$

17th Mar, 2021 (shift 1)

$$A^2 = \begin{bmatrix} \sin^2 \alpha & 0 \\ 0 & \sin^2 \alpha \end{bmatrix}$$

$$\left| A^2 - \frac{1}{2}I \right| = \begin{vmatrix} \sin^2 \alpha - \frac{1}{2} & 0 \\ 0 & \sin^2 \alpha - \frac{1}{2} \end{vmatrix} = 0$$

$$\left( \sin^2 \alpha - \frac{1}{2} \right)^2 = 0 \Rightarrow \boxed{\sin^2 \alpha = \frac{1}{2}}$$



Let A be a  $2 \times 2$  matrix with  $\det(A) = -1$  and  $\det((A + I)(\text{Adj}(A) + I)) = 4$ . Then the sum of the diagonal elements of A can be :

(A)  $-1$

~~(B)  $2$~~

(C)  $1$

$a+d = 2/-2$

(D)  $-\sqrt{2}$

$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \boxed{ad - bc = -1}$

$| (A+I)(\text{adj} A + I) | = 4$

$|A+I| |\text{adj} A + I| = 4$

$(a+d)(a+d) = 4$

$(a+d)^2 = 4$

$a+d = 2 \text{ or } -2$

$|A+I| = \begin{bmatrix} a+1 & b \\ c & d+1 \end{bmatrix}$

$= (a+1)(d+1) - bc$

$= \cancel{ad} + a + d + \cancel{1} - \cancel{bc}$

$= a+d$

*JEE Main 2022*

2x2

$$(a+d)^2 = 4$$

$$\text{adj } A = \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

$$\begin{aligned} |\text{adj } A + I| &= \begin{vmatrix} d+1 & -b \\ -c & a+1 \end{vmatrix} \\ &= (d+1)(a+1) - bc \\ &= a+d+1+ad-bc \\ &= a+d \end{aligned}$$

$$\begin{aligned} A \text{ adj } A &= |A| I \\ &= \underline{-I} \end{aligned}$$

$$|\cancel{A \text{ adj } A} + A + \text{adj } A + \cancel{I}| = 4$$

$$|A + \text{adj } A| = 4$$

$$\left| \begin{bmatrix} a & b \\ c & d \end{bmatrix} + \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \right| = 4$$

$$\begin{vmatrix} a+d & 0 \\ 0 & a+d \end{vmatrix} = 4$$

Domain

Let  $|M|$  denote the determinant of a square matrix  $M$ . Let  $g: \left[0, \frac{\pi}{2}\right] \rightarrow \mathbb{R}$  be the function defined by

AC

$$g(\theta) = \sqrt{f(\theta) - 1} + \sqrt{f\left(\frac{\pi}{2} - \theta\right) - 1}$$

where

$$f(\theta) = \frac{1}{2} \begin{vmatrix} 1 & \sin \theta & 1 \\ -\sin \theta & 1 & \sin \theta \\ -1 & -\sin \theta & 1 \end{vmatrix} + \begin{vmatrix} \sin \pi & \cos\left(\theta + \frac{\pi}{4}\right) & \tan\left(\theta - \frac{\pi}{4}\right) \\ \sin\left(\theta - \frac{\pi}{4}\right) & -\cos\frac{\pi}{2} & \log_e\left(\frac{4}{\pi}\right) \\ \cot\left(\theta + \frac{\pi}{4}\right) & \log_e\left(\frac{\pi}{4}\right) & \tan \pi \end{vmatrix}.$$

Let  $p(x)$  be a quadratic polynomial whose roots are the maximum and minimum values of the function  $g(\theta)$ , and  $p(2) = 2 - \sqrt{2}$ . Then, which of the following is/are TRUE?

(A)  $p\left(\frac{3+\sqrt{2}}{4}\right) < 0$

$$\frac{3+1.4}{4} = 1.1$$

(B)  $p\left(\frac{1+3\sqrt{2}}{4}\right) > 0$

$$\frac{1+3(1.4)}{4} = 1.3$$

(C)  $p\left(\frac{5\sqrt{2}-1}{4}\right) > 0$

$$\frac{5(1.4)-1}{4} = 1.5$$

(D)  $p\left(\frac{5-\sqrt{2}}{4}\right) < 0$

$$\frac{5-1.4}{4} = 0.9$$

$$\begin{aligned} & \cos\left(\theta + \frac{\pi}{4}\right) \\ &= \sin\left(\frac{\pi}{2} - \theta - \frac{\pi}{4}\right) \\ &= \sin\left(\frac{\pi}{4} - \theta\right) \\ &= -\sin\left(\theta - \frac{\pi}{4}\right) \end{aligned}$$

JEE Adv. 2022

$$\begin{vmatrix} 0 & k & - \\ -k & 0 & - \\ - & - & 0 \end{vmatrix}$$

Skew sym + odd order

$$|\mathbb{A}| = 0$$

$$f(\theta) = \frac{1}{2} \begin{vmatrix} 2 & \sin \theta & 1 \\ 0 & 1 & \sin \theta \\ 0 & -\sin \theta & 1 \end{vmatrix}$$

$$= \frac{1}{2} \cancel{2} (1 + \sin^2 \theta)$$

$$\boxed{f(\theta) = 1 + \sin^2 \theta}$$

$$g(\theta) = \sqrt{f(\theta) - 1} + \sqrt{f(\frac{\pi}{2} - \theta) - 1}$$

$$= \sqrt{1 + \sin^2 \theta - 1} + \sqrt{1 + \cos^2 \theta - 1}$$

$$= |\sin \theta| + |\cos \theta| = \underline{\sin \theta + \cos \theta}$$

$$g(0) = 1 \quad g(\frac{\pi}{4}) = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} = \sqrt{2}$$

$$g(\frac{\pi}{2}) = 1$$

$$\boxed{y(\theta) = \sin \theta + \cos \theta} \quad \begin{cases} \text{max} = \sqrt{2} \\ \text{min} = 1 \end{cases}$$


$$p(x) = K(x - \sqrt{2})(x - 1)$$

$$\cancel{2 - \sqrt{2}} = K(\cancel{2 - \sqrt{2}})(2 - 1)$$

$$\therefore \boxed{K = 1}$$

$$p(x) = (x - \sqrt{2})(x - 1)$$

GRAPH





$$p(x) = (x - 1.414)(x - 1)$$

$$p(1.1) = (1.1 - 1.4)(1.1 - 1) = \ominus$$

$$p(1.3) = (1.3 - 1.4)(1.3 - 1) = \ominus$$

$$p(1.5) = (1.5 - 1.4)(1.5 - 1) = \oplus$$

$$p(0.9) = (0.9 - 1.4)(0.9 - 1) = \oplus$$





# Special Determinants



# Special Determinants



$$\star \begin{vmatrix} 1 & x & x^2 \\ 1 & y & y^2 \\ 1 & z & z^2 \end{vmatrix} = (x - y)(y - z)(z - x)$$

$$\star \begin{vmatrix} 1 & x & x^3 \\ 1 & y & y^3 \\ 1 & z & z^3 \end{vmatrix} = (x - y)(y - z)(z - x)(x + y + z)$$

$$\star \star \star \begin{vmatrix} a & b & c \\ b & c & a \\ c & a & b \end{vmatrix} = -(a^3 + b^3 + c^3 - 3abc)$$




# Special Determinants



$$\begin{vmatrix} 1 & x & x^2 \\ 1 & y & y^2 \\ 1 & z & z^2 \end{vmatrix} = \frac{(x-y)(y-z)(z-x)}{1}$$

$$R_2 \rightarrow R_2 - R_1$$

$$R_3 \rightarrow R_3 - R_1$$



$$\begin{vmatrix} 1 & x & x^2 \\ 0 & y-x & (y-x)(y+x) \\ 0 & z-x & (z-x)(z+x) \end{vmatrix}$$

$$= (y-x)(z-x)$$

$$\begin{vmatrix} 1 & x & x^2 \\ 0 & 1 & y+x \\ 0 & 1 & z+x \end{vmatrix}$$

$$= (y-x)(z-x)(z-y)$$

$$= \underline{(x-y)(y-z)(z-x)}$$





Prove that  $\begin{vmatrix} \underline{b+c} & c & b \\ c & \underline{c+a} & a \\ b & a & \underline{a+b} \end{vmatrix} = \underline{4abc}$

✓ 1 1 1

✓ 0 0

$$R_1 \rightarrow R_1 - (R_2 + R_3)$$

$$\begin{vmatrix} \boxed{0} & -2a & -2a \\ c & c+a & a \\ b & a & a+b \end{vmatrix} = -2a \begin{vmatrix} 0 & 1 & 1 \\ c & c+a & a \\ b & a & a+b \end{vmatrix}$$
$$= -2a \begin{vmatrix} \text{---} & \text{---} & \text{---} \\ c & c & a \\ b & -b & a+b \end{vmatrix}$$
$$= -2a(-2bc)$$
$$= \underline{4abc}$$





Show that 
$$\begin{vmatrix} \mathbf{b + c} & \mathbf{a + b} & \mathbf{a} \\ \mathbf{c + a} & \mathbf{b + c} & \mathbf{b} \\ \mathbf{a + b} & \mathbf{c + a} & \mathbf{c} \end{vmatrix} = \mathbf{a^3 + b^3 + c^3 - 3abc}.$$

*H w*









Show that 
$$\begin{vmatrix} a-b-c & 2a & 2a \\ 2b & b-c-a & 2b \\ 2c & 2c & c-a-b \end{vmatrix} = (a+b+c)^3.$$

H w







Show that

$$\begin{vmatrix} x & x^2 & yz \\ y & y^2 & zx \\ z & z^2 & xy \end{vmatrix} = (x - y)(y - z)(z - x)(xy + yz + zx)$$

Concept

#chalaki



$$\begin{array}{ccc|ccc} xx & \rightarrow & x^2 & x^3 & yz & \\ xy & \rightarrow & y^2 & y^3 & zx & \\ xz & \rightarrow & z^2 & z^3 & xy & \end{array} \quad = \quad \begin{vmatrix} x^2 & x^3 & 1 \\ y^2 & y^3 & 1 \\ z^2 & z^3 & 1 \end{vmatrix}$$

$\downarrow$   
 $xyz$

$$\left| \begin{array}{cc|c} x^2 & x^3 & 1 \\ \hline y^2 - x^2 & y^3 - x^3 & 0 \\ \hline z^2 - x^2 & z^3 - x^3 & 0 \end{array} \right|$$

$$(y-x)(z-x) \left| \begin{array}{cc|c} x^2 & x^3 & 1 \\ y+x & y^2+x^2+xy & 0 \\ z+x & z^2+x^2+xz & 0 \end{array} \right|$$





Show that  $\begin{vmatrix} a^2 + 1 & ab & ac \\ ab & b^2 + 1 & bc \\ ac & bc & c^2 + 1 \end{vmatrix} = \underline{1 + a^2 + b^2 + c^2}.$

#chalaki

$$\begin{array}{l} a \rightarrow \\ b \rightarrow \\ c \rightarrow \end{array} \begin{vmatrix} (a^2 + 1) & ab & ac \\ ab^2 & (b^2 + 1) & bc \\ ac^2 & bc^2 & (c^2 + 1) \end{vmatrix} = \begin{vmatrix} a^2 + 1 & a^2 & a^2 \\ b^2 & b^2 + 1 & b^2 \\ c^2 & c^2 & c^2 + 1 \end{vmatrix}$$

$\downarrow \quad \downarrow \quad \downarrow$   
 $a \quad b \quad c$

$$R_1 \rightarrow R_1 + R_2 + R_3$$

$$(1 + a^2 + b^2 + c^2) \begin{vmatrix} 1 & 1 & 1 \\ b^2 & b^2 + 1 & b^2 \\ c^2 & c^2 & c^2 + 1 \end{vmatrix}$$

$$(1+a^2+b^2+c^2) \begin{vmatrix} 1 & 0 & 0 \\ b^2 & 1 & 0 \\ c^2 & 0 & 1 \end{vmatrix}$$

$$= \underline{1+a^2+b^2+c^2}$$







Which of the following values of  $\alpha$  satisfy the equation

$$\begin{vmatrix} (1+\alpha)^2 & (1+2\alpha)^2 & (1+3\alpha)^2 \\ (2+\alpha)^2 & (2+2\alpha)^2 & (2+3\alpha)^2 \\ (3+\alpha)^2 & (3+2\alpha)^2 & (3+3\alpha)^2 \end{vmatrix} = -648\alpha ?$$

(JEE Adv. 2015)

A. -4

B. 9

C. -9

D. 4

$$R_2 \rightarrow R_2 - R_1$$

$$R_3 \rightarrow R_3 - R_2$$

$$\begin{vmatrix} (1+\alpha)^2 & (1+2\alpha)^2 & (1+3\alpha)^2 \\ 2\alpha+3 & 4\alpha+3 & 6\alpha+3 \\ 2\alpha+5 & 4\alpha+5 & 6\alpha+5 \end{vmatrix} = -648\alpha$$

$$R_3 \rightarrow R_3 - R_2$$

$$2 \begin{vmatrix} \overbrace{(1+\alpha)^2} & \overbrace{(1+2\alpha)^2} & \overbrace{(1+3\alpha)^2} \\ \underline{2\alpha+3} & \underline{4\alpha+3} & 6\alpha+3 \\ | & | & | \end{vmatrix} = -648\alpha$$

$$2 \begin{vmatrix} \overbrace{(1+\alpha)^2} & 3\alpha^2+2\alpha & 5\alpha^2+2\alpha \\ 2\alpha+3 & 2\alpha & 2\alpha \\ || & 0 & 0 \end{vmatrix} = -648\alpha$$

$\cancel{2} \left( \cancel{+4} \alpha^3 \right) = \cancel{+648} \alpha$

$$\alpha^3 - 81\alpha = 0$$

$$\alpha(\alpha-9)(\alpha+9) = 0$$

$$\alpha = 0, 9, -9$$





If  $a^2 + b^2 + c^2 = -2$  and

$$a^2 + b^2 + c^2 + 2 = 0$$

$$f(x) = \begin{vmatrix} \frac{1+a^2x}{(1+a^2)x} & \frac{(1+b^2)x}{1+b^2x} & \frac{(1+c^2)x}{(1+c^2)x} \\ (1+a^2)x & 1+b^2x & (1+c^2)x \\ (1+a^2)x & (1+b^2)x & 1+c^2x \end{vmatrix}$$

Hint

then  $f(x)$  is a polynomial degree

(JEE Adv. 2005)

A. 1

B. 0

C. 3

☒ D. 2

$$C_1 \rightarrow C_1 + C_2 + C_3$$

$$1 + (a^2 + b^2 + c^2 + 2)x$$

$$1 + (a^2 + b^2 + c^2 + 2)x$$

$$1 + (a^2 + b^2 + c^2 + 2)x$$

—

—

—

—

—

—

$$f(x) = \begin{vmatrix} 1 & \underline{(1+b^2)x} & \underline{(1+c^2)x} \\ \textcircled{1} & \underline{1+b^2x} & \underline{(1+c^2)x} \\ \textcircled{1} & \underline{(1+b^2)x} & \underline{1+c^2x} \end{vmatrix}$$

$$R_2 \rightarrow R_2 - R_1$$

$$R_3 \rightarrow R_3 - R_1$$

$$f(x) = \begin{vmatrix} \textcircled{1} & (1+b^2)x & (1+c^2)x \\ 0 & 1-x & 0 \\ 0 & 0 & 1-x \end{vmatrix}$$

$$f(x) = (1-x)^2 = 1 - 2x + x^2$$





Find **values of**

$$\begin{vmatrix} \sin \theta & \cos \theta & \sin 2\theta \\ \sin\left(\theta + \frac{2\pi}{3}\right) & \cos\left(\theta + \frac{2\pi}{3}\right) & \sin\left(2\theta + \frac{4\pi}{3}\right) \\ \sin\left(\theta - \frac{2\pi}{3}\right) & \cos\left(\theta - \frac{2\pi}{3}\right) & \sin\left(2\theta - \frac{4\pi}{3}\right) \end{vmatrix}$$

$$R_3 \rightarrow R_3 + R_2$$

$$\begin{vmatrix} \sin \theta & \cos \theta & \sin 2\theta \\ \sin\left(\theta + \frac{2\pi}{3}\right) & - & - \\ \cancel{2 \sin \theta \left(\frac{-1}{2}\right)} & \cancel{2 \cos \theta \left(\frac{-1}{2}\right)} & \cancel{2 \sin 2\theta \left(\frac{-1}{2}\right)} \end{vmatrix} = 0$$

(2000 - 3 marks)



$$\cos \frac{2\pi}{3} = \cos\left(\pi - \frac{\pi}{3}\right) = -\cos \frac{\pi}{3} = -\frac{1}{2}$$

$$\cos\left(\frac{4\pi}{3}\right) = \cos\left(\pi + \frac{\pi}{3}\right) = \left(-\frac{1}{2}\right)$$





# ✓ Method to Solve System of Linear Equations



# System Linear Equations



**1**

**Determinant Method (Cramer's Rule)**

**2**

**Matrix Method (Gauss- Jordan Method)**



# Cramer's Rule



# Cramer's Rule

Const  $\rightarrow$  Right Side

$$\left\{ \begin{array}{l} a_1x + b_1y + c_1z = d_1 \\ a_2x + b_2y + c_2z = d_2 \\ a_3x + b_3y + c_3z = d_3 \end{array} \right\} \text{system of L.E}$$

$$x = \frac{D_1}{D}$$

$$y = \frac{D_2}{D}$$

$$z = \frac{D_3}{D}$$

$$D = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

$$D = 0$$

$\swarrow \searrow$   
 $N_0 \quad \infty$

$$D \neq 0$$

$$\left. \begin{array}{l} x = \\ y = \\ z = \end{array} \right\} \text{Unique Sol}^n$$

$$D_1 = \begin{vmatrix} d_1 & b_1 & c_1 \\ d_2 & b_2 & c_2 \\ d_3 & b_3 & c_3 \end{vmatrix}$$

$$D_2 = \begin{vmatrix} a_1 & d_1 & c_1 \\ a_2 & d_2 & c_2 \\ a_3 & d_3 & c_3 \end{vmatrix}$$

$$D_3 = \begin{vmatrix} a_1 & b_1 & d_1 \\ a_2 & b_2 & d_2 \\ a_3 & b_3 & d_3 \end{vmatrix}$$

# Cramer's Rule

$$x = \frac{D_1}{D} \quad \frac{0}{0}$$

$\neq$

If  $D \neq 0$   
Unique solution

If  $D = 0$

$D \neq 0$   
Unique

$D = 0$

$D_1, D_2, D_3$   
 $\neq 0$   
No soln

$D_1 = D_2 = D_3 = 0$   
~~No (parallel)~~  
 $\infty$   
( $\infty$ )

If at least one of  
 $D_1, D_2, D_3 \neq 0$   
No solution

$D_1, D_2, D_3$  Koi ek  $\neq 0$   
No soln

If  $D_1 = D_2 = D_3 = 0$   
Infinite Solution Or  
No Solution

planes are parallel




## Important terms



- i. **Consistent:** solution exists (unique or infinite solution)
- ii. **Inconsistent:** solution does not exist (No solution)
- iii. **Homogeneous equations:** constant terms are zero
- iv. **Trivial solution:** all variables = zero i.e.,  $x = 0, y = 0, z = 0$ .

$$x = 0 \quad y = 0 \quad z = 0$$

$$\boxed{\text{all Var} = 0}$$





# Homogeneous Linear Equations



$$\begin{cases} a_1x + b_1y + c_1z = 0 \\ a_2x + b_2y + c_2z = 0 \\ a_3x + b_3y + c_3z = 0 \end{cases}$$

*Homo Linear Eqn.*

$$D_1 = D_2 = D_3 = 0$$

$$D_1 = 0 = D_2 = D_3$$

$$D = 0 \text{ or } D \neq 0$$


$$D = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

$$D_1 = \begin{vmatrix} 0 & - & - \\ 0 & - & - \\ 0 & - & - \end{vmatrix} = 0$$

$$D_2 = \begin{vmatrix} 0 & & \\ 0 & & \\ 0 & & \end{vmatrix} = 0$$

$$D_3 = \begin{vmatrix} 0 & \\ 0 & \\ 0 & \end{vmatrix} = 0$$







# Gauss-Jordan Method



# Matrix Method (Gauss-Jordan Method)

$$x + y + z = 6 \checkmark$$

$$x - y + z = 2 \checkmark$$

$$2x + y - z = 1$$

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \\ 2 & 1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 6 \\ 2 \\ 1 \end{bmatrix}$$

$$AX = B$$

$$\tilde{A}^{-1} \tilde{A} X = \tilde{A}^{-1} B$$

$$X = \tilde{A}^{-1} B$$

$$x = \frac{D_1}{D} \quad y = \frac{D_2}{D} \quad z = \frac{D_3}{D}$$

$$X = \frac{(\text{adj } A) B}{|A|}$$

$$|A| \neq 0$$

# Unique

$$|A| = 0$$

$$(\text{adj } A) \cdot B = 0$$

(infinite)


$$(\text{adj } A) B \neq 0$$

(No Sol<sup>n</sup>)

$\frac{0}{0} \quad \infty$



# Matrix Method (Gauss-Jordan Method)





# Questions



For what values of  $p$  and  $q$  the system of equations has

$$2x + py + 6z = 8$$

$$x + 2y + qz = 5$$

$$x + y + 3z = 4$$

- i. Unique solution      ii. No solution      iii. Infinite solutions



i) Unique Sol<sup>n</sup>

$$D \neq 0$$

$$(p-2)(q-3) \neq 0$$

$$\boxed{p \neq 2 \text{ and } q \neq 3}$$

ii) No Sol<sup>n</sup>  $D=0$   $(D_1, D_2, D_3)$  Koi ek  $\neq 0$

$$\boxed{p \neq 2, q = 3}$$

iii)  $\infty$  Sol<sup>n</sup>  $D = D_1 = D_2 = D_3 = 0$

$$\boxed{p = 2; q \in \mathbb{R}}$$

$$D = \begin{vmatrix} 2 & p & 6 \\ 1 & 2 & q \\ 1 & 1 & 3 \end{vmatrix} = (p-2)(q-3)$$

$$D_1 = \begin{vmatrix} 8 & p & 6 \\ 5 & 2 & q \\ 4 & 1 & 3 \end{vmatrix} = (p-2)(4q-15)$$

$$D_2 = \begin{vmatrix} 2 & 8 & 6 \\ 1 & 5 & q \\ 1 & 4 & 3 \end{vmatrix} = 0$$

$$D_3 = \begin{vmatrix} 2 & p & 8 \\ 1 & 2 & 5 \\ 1 & 1 & 4 \end{vmatrix} = (p-2)$$





The system of equations

$$kx + y + z = 1$$

$$x + ky + z = k$$

$$x + y + zk = k^2$$

has **no solution**, if  $k$  is equal to -

$$D = 0$$

$(D_1, D_2, D_3)$  at least one  $\neq 0$

17th March 2021, Shift 1

~~A.~~ 0

B. 1

~~C.~~ -1

☒ D. -2

$$D = \begin{vmatrix} k & 1 & 1 \\ 1 & k & 1 \\ 1 & 1 & k \end{vmatrix} = (k+2)(k-1)^2$$

$$D_1 = \begin{vmatrix} 1 & 1 & 1 \\ k & k & 1 \\ k^2 & 1 & k \end{vmatrix} = -(k-1)^2(k+1)$$

$$D=0 \quad k = -2/1$$

✓ X

if  $k \neq -2$

$$\begin{matrix} D=0 \\ D_1 \neq 0 \end{matrix}$$

$k=1$





Let the system of linear equations

$$4x + \lambda y + 2z = 0$$

$$2x - y + z = 0$$

$$\mu x + 2y + 3z = 0, \lambda, \mu \in \mathbb{R}$$

Homo + Non-trivial

$$D = 0$$

has **non-trivial** solution, then which of the following is true?

$$D = \begin{vmatrix} 4 & \lambda & 2 \\ 2 & -1 & 1 \\ \mu & 2 & 3 \end{vmatrix} = \underline{(\lambda + 2)(\mu - 6)}$$

18th March 2021, Shift 2

☒ A.  $\mu = 6, \lambda \in \mathbb{R}$

☐ B.  $\lambda = 2, \mu \in \mathbb{R}$

☐ C.  $\lambda = 3, \mu \in \mathbb{R}$

☐ D.  $\mu = -6, \lambda \in \mathbb{R}$



For the system of linear equations

$$x - 2y = 1$$

$$x - y + kz = -2$$

$$ky + 4z = 6, k \in \mathbb{R}$$

24th February 2021, Shift 2

Consider the following statements:

1. The system has unique solution if  $k \neq 2, k \neq -2$
- ~~2. The system has unique solution if  $k = -2$~~
- ~~3. The system has unique solution if  $k = 2$~~
4. The system has no solution if  $k = 2$
- ~~5. The system has infinite number of solutions if  $k \neq -2$~~

Which of the following statements are correct?

- ~~A. 2 & 5 only~~
- ~~B. 3 & 4 only~~
- ☒ C. 1 & 4 only
- ~~D. 1 & 5 only~~

$$k = -2 \rightarrow \infty \text{ soln}$$



$$D = \begin{vmatrix} 1 & -2 & 0 \\ 1 & -1 & k \\ 0 & k & 4 \end{vmatrix} = (2 - k)(2 + k)$$

$$D_1 = \begin{vmatrix} 1 & -2 & 0 \\ -2 & -1 & k \\ 6 & k & 4 \end{vmatrix} = -(k + 2)(k + 10)$$

$$D_2 = \begin{vmatrix} 1 & 1 & 0 \\ 1 & -2 & k \\ 0 & 6 & 4 \end{vmatrix} = -6(k + 2)$$

$$D_3 = \begin{vmatrix} 1 & -2 & 1 \\ 1 & -1 & -2 \\ 0 & k & 6 \end{vmatrix} = 3(k + 2)$$

if  $k = 2$

$$\left. \begin{array}{l} D = 0 \\ D_3 \neq 0 \end{array} \right\}$$

Unique Sol<sup>n</sup>

$$D \neq 0$$

$$\boxed{k \neq 2, -2}$$

$\infty$  Sol<sup>n</sup>

$$\boxed{k = -2}$$



If the system of equations

$$kx + y + 2z = 1$$

$$3x - y - 2z = 2$$

$$-2x - 2y - 4z = 3$$

has infinitely many solutions, then  $k$  is equal to -

$$D = \begin{vmatrix} k & 1 & 2 \\ 3 & -1 & -2 \\ -2 & -2 & -4 \end{vmatrix} = 0$$

$$D_1 = \begin{vmatrix} 1 & 1 & 2 \\ 2 & -1 & -2 \\ 3 & -2 & -4 \end{vmatrix} = 0$$

$$D_2 = \begin{vmatrix} k & 1 & 2 \\ 3 & 2 & -2 \\ -2 & 3 & -4 \end{vmatrix} = -2(k - 21)$$

$$D_3 = \begin{vmatrix} k & 1 & 1 \\ 3 & -1 & 2 \\ -2 & -2 & 3 \end{vmatrix} = (k - 21)$$

25th February 2021, Shift 1

$$K = 21$$



The following system of linear equations

$$2x + 3y + 2z = 9$$

$$3x + 2y + 2z = 9$$

$$x - y + 4z = 8$$

*25th February 2021, Shift 2*



does not have any solution



has a unique solution



has a solution  $(\alpha, \beta, \gamma)$  satisfying  $\alpha + \beta^2 + \gamma^3 = 12$

$$0 + 1^2 + 2^3 \neq 12$$



has infinitely many solutions



$$D = \begin{vmatrix} 2 & 3 & 2 \\ 3 & 2 & 2 \\ 1 & -1 & 4 \end{vmatrix} = -20$$

$$D_1 = \begin{vmatrix} \color{red}{9} & 3 & 2 \\ \color{red}{9} & 2 & 2 \\ \color{red}{8} & -1 & 4 \end{vmatrix} = 0$$

$$D_2 = \begin{vmatrix} 2 & \color{red}{9} & 2 \\ 3 & \color{red}{9} & 2 \\ 1 & \color{red}{8} & 4 \end{vmatrix} = -20$$

$$D_3 = \begin{vmatrix} 2 & 3 & \color{red}{9} \\ 3 & 2 & \color{red}{9} \\ 1 & -1 & \color{red}{8} \end{vmatrix} = -40$$

$$D \neq 0$$

Unique

$$x = \frac{D_1}{D} = 0$$

$$y = \frac{D_2}{D} = 1$$

$$z = \frac{D_3}{D} = 2$$

(0, 1, 2)  
Sol<sup>n</sup>



If the following system of linear equations

$$2x + y + z = 5$$

$$x - y + z = 3$$

$$x + y + az = b$$

has no solution, then :

$$\boxed{D=0} \quad D_1, D_2, D_3 \neq 0$$

$$a = \frac{1}{3}$$

31 Aug 2021, Shift 1

$$a = \frac{1}{3} \quad b \neq \frac{7}{3}$$

$$\boxed{D=0 \quad D_3 \neq 0}$$

No Sol<sup>n</sup>

☒ A.  $a = -\frac{1}{3}, b \neq \frac{7}{3}$

☒ B.  $a \neq \frac{1}{3}, b = \frac{7}{3}$

☒ C.  $a \neq -\frac{1}{3}, b = \frac{7}{3}$

☒ D.  $a = \frac{1}{3}, b \neq \frac{7}{3}$

$$D = \begin{vmatrix} 2 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & a \end{vmatrix} = 1 - 3a$$

$$D_3 = \begin{vmatrix} 2 & 1 & 5 \\ 1 & -1 & 3 \\ 1 & 1 & b \end{vmatrix} = 7 - 3b$$



The value of a and b, for which the system of equations

$$2x + 3y + 6z = 8$$

$$x + 2y + az = 5$$

$$3x + 5y + 9z = b$$

has no solution, are :

*25 July 2021, Shift 1*

$$D=0$$

☒ A.  $a = 3, b \neq 13$

☐ B.  $a \neq 3, b \neq 13$

☐ C.  $a \neq 3, b = 3$

☐ D.  $a = 3, b = 13$

$$D = \begin{vmatrix} 2 & 3 & 6 \\ 1 & 2 & a \\ 3 & 5 & 9 \end{vmatrix} = 3 - a = 0$$

$$D_3 = \begin{vmatrix} 2 & 3 & 8 \\ 1 & 2 & 5 \\ 3 & 5 & b \end{vmatrix} = b - 13 \neq 0$$





The value of  $\lambda$  and  $\mu$  such that the system of equations  $x + y + z = 6$ ,  $3x + 5y + 5z = 26$ ,  $x + 2y + \lambda z = \mu$  has no solutions, are :

A.  $\lambda = 3, \mu = 5$

B.  $\lambda = 3, \mu \neq 10$

C.  $\lambda \neq 2, \mu = 10$

☒ D.  $\lambda = 2, \mu \neq 10$

*22 July 2021, shift 1*

$$D = \begin{vmatrix} 1 & 1 & 1 \\ 3 & 5 & 5 \\ 1 & 2 & \lambda \end{vmatrix} = 2\lambda - 4$$

$$D = 0$$
$$\boxed{\lambda = 2}$$



The value of  $k \in \mathbb{R}$ , for which the following system of linear equations

$$3x - y + 4z = 3$$

$$x + 2y - 3z = -2$$

$$6x + 5y + kz = -3$$

Has infinitely many solutions is :

*20 July 2021, shift 2*

A. 3

☒ B. -5

C. 5

D. -3

$$\underline{D} = D_1 = D_2 = D_3 = 0$$

$$D = \begin{vmatrix} 3 & -1 & 4 \\ 1 & 2 & -3 \\ 6 & 5 & k \end{vmatrix} = 7k + 35 = 0$$

$$k = -5$$



# Cramer's Rule (JEE Main 2022)



If the system of equations

$$x + y + z = 6$$

$$2x + 5y + \alpha z = \beta$$

$$x + 2y + 3z = 14$$

Has infinitely many solutions, then  $\alpha + \beta$  is equal to :

*JEE Main 2022*

A. 8

B. 36

☒ C. 44

D. 48

$$D = \begin{vmatrix} 1 & 1 & 1 \\ 2 & 5 & \alpha \\ 1 & 2 & 3 \end{vmatrix} = 8 - \alpha = 0$$

$$D_3 = \begin{vmatrix} 1 & 1 & 6 \\ 2 & 5 & \beta \\ 1 & 2 & 14 \end{vmatrix} = 36 - \beta = 0$$

$$\begin{array}{r} \alpha = 8 \\ + \beta = 36 \\ \hline \alpha + \beta = 44 \end{array}$$



The number of values of  $\alpha$  for which the system of equations :

$$x + y + z = \alpha$$

$$\alpha x + 2\alpha y + 3z = -1$$

$$x + 3\alpha y + 5z = 4$$

*JEE Main 2022*

Is inconsistent, is

$D = 0$

*No soln*

A.

0

$$\begin{vmatrix} 1 & 1 & 1 \\ \alpha & 2\alpha & 3 \\ 1 & 3\alpha & 5 \end{vmatrix} = 3(\alpha - 1)^2 = 0$$

☒ B.

1

C.

2

D.

3

$$\begin{vmatrix} \alpha & 1 & 1 \\ -1 & 2\alpha & 3 \\ 4 & 3\alpha & 5 \end{vmatrix} = \alpha^2 - 11\alpha + 7 \neq 0$$



Let the system of linear equations

$$x + y + \alpha z = 2$$

$$3x + y + z = 4$$

$$x + 2z = 1$$

have a unique solution  $(x^*, y^*, z^*)$ . If  $(\alpha, x^*)$ ,  $(y^*, \alpha)$  and  $(x^*, -y^*)$  are collinear points, then the sum of absolute values of all possible values of  $\alpha$  is:

*JEE Main 2022*

A.

4

B.

3

C.

2

D.

1

$$|1| + |-1|$$

$$\Rightarrow (2)$$

$(\alpha, 1)$ ,  $(1, \alpha)$  and  $(1, -1)$   
Collinear

$$\begin{vmatrix} \alpha & 1 & 1 \\ 1 & \alpha & 1 \\ 1 & -1 & 1 \end{vmatrix} = 0$$

$$\alpha(\alpha+1) - 1(0) + 1(-1-\alpha) = 0$$

$$\alpha^2 + \cancel{\alpha} - 1 - \cancel{\alpha} = 0$$

$$\boxed{\alpha = \pm 1}$$



$$\Delta = \begin{vmatrix} 1 & 1 & \alpha \\ 3 & 1 & 1 \\ 1 & 0 & 2 \end{vmatrix} = -(\alpha + 3)$$

$$\Delta_1 = \begin{vmatrix} 2 & 1 & \alpha \\ 4 & 1 & 1 \\ 1 & 0 & 2 \end{vmatrix} = -(3 + \alpha)$$

$$\Delta_2 = \begin{vmatrix} 1 & 2 & \alpha \\ 3 & 4 & 1 \\ 1 & 1 & 2 \end{vmatrix} = -(\alpha + 3)$$

$$\Delta_3 = \begin{vmatrix} 1 & 1 & 2 \\ 3 & 1 & 4 \\ 1 & 0 & 1 \end{vmatrix} = 0$$

$$x^* = 1 \quad y^* = 1 \quad z^* = 0$$



The ordered pair  $(a, b)$ , for which the system of linear equations

$$3x - 2y + z = b$$

$$5x - 8y + 9z = 3$$

$$2x + y + az = -1$$

Has no solution, is :

*JEE Main 2022*

A.  $\left(3, \frac{1}{3}\right)$

$$D = \begin{vmatrix} 3 & -2 & 1 \\ 5 & -8 & 9 \\ 2 & 1 & a \end{vmatrix} = -14(a + 3) = 0$$

B.  $\left(-3, \frac{1}{3}\right)$

☒ C.  $\left(-3, -\frac{1}{3}\right)$

D.  $\left(3, -\frac{1}{3}\right)$

$$D_3 = \begin{vmatrix} 3 & -2 & b \\ 5 & -8 & 3 \\ 2 & 1 & -1 \end{vmatrix} = 7(3b - 1) \neq 0$$

$$a = -3$$

$$b \neq \frac{1}{3}$$





If the system of equations  $\alpha x + y + z = 5$ ,  $x + 2y + 3z = 4$ ,  $x + 3y + 5z = \beta$ , has **infinitely** many solutions, then the ordered pair  $(\alpha, \beta)$  is equal to:

*JEE Main 2022*

A. (1, -3)

B. (-1, 3)

☒ C. (1, 3)

D. (-1, -3)

$$D = \begin{vmatrix} \alpha & 1 & 1 \\ 1 & 2 & 3 \\ 1 & 3 & 5 \end{vmatrix} = \alpha - 1 = 0$$

$$D_1 = \begin{vmatrix} 5 & 1 & 1 \\ 4 & 2 & 3 \\ \beta & 3 & 5 \end{vmatrix} = \beta - 3 = 0$$

$$\underline{D = D_1 = D_2 = D_3 = 0}$$



Let the system of linear equations  $x + 2y + z = 2$ ,  $\alpha x + 3y - z = \alpha$ ,  $-\alpha x + y + 2z = -\alpha$  be inconsistent. Then  $\alpha$  is equal to :

No Sol<sup>n</sup>

*JEE Main 2022*

A.  $5/2$

B.  $-5/2$

C.  $7/2$

☒ D.  $-7/2$

$$D = \begin{vmatrix} 1 & 2 & 1 \\ \alpha & 3 & -1 \\ -\alpha & 1 & 2 \end{vmatrix} = 7 + 2\alpha = 0$$

$$D_1 = \begin{vmatrix} 2 & 2 & 1 \\ \alpha & 3 & -1 \\ -\alpha & 1 & 2 \end{vmatrix} = 14 + 2\alpha \neq 0$$

$\alpha = -7/2$



If the system of linear equations

$$2x + y - z = 7$$

$$x - 3y + 2z = 1$$

$$x + 4y + \delta z = k, \text{ where } \delta, k \in \mathbb{R}$$

has infinitely many solutions, then  $\delta + k$  is equal to :

*JEE Main 2022*

A.

-3

B.

3

C.

6

D.

9

$$D = \begin{vmatrix} 2 & 1 & -1 \\ 1 & -3 & 2 \\ 1 & 4 & \delta \end{vmatrix} = -7(\delta + 3) = 0$$

$$D_3 = \begin{vmatrix} 2 & 1 & 7 \\ 1 & -3 & 1 \\ 1 & 4 & k \end{vmatrix} = -7(k - 6) = 0$$

$$\delta = -3$$

$$k = 6$$

$$\underline{\delta + k = 3}$$

Let  $p, q, r$  be nonzero real numbers that are, respectively, the  $10^{th}$ ,  $100^{th}$  and  $1000^{th}$  terms of a harmonic progression. Consider the system of linear equations

$$\frac{1}{p}, \frac{1}{q}, \frac{1}{r} \rightarrow AP$$

$$\begin{aligned} &\checkmark x + y + z = 1 \\ &10x + 100y + 1000z = 0 \\ &\underline{qr x + pr y + pq z = 0} \end{aligned}$$

$$pqr$$

$$\begin{aligned} \frac{1}{p} &= A + 9D \\ \frac{1}{q} &= A + 99D \\ \frac{1}{r} &= A + 999D \end{aligned}$$

**JEE Adv. 2022**

$$\frac{p}{q} = \frac{A + 99D}{A + 9D}$$

### List-I

### List-II

(I) If  $\frac{q}{r} = 10$ , then the system of linear equations has

(P)  $x = 0$ ,  $y = \frac{10}{9}$ ,  $z = -\frac{1}{9}$  as a solution

(II) If  $\frac{p}{r} \neq 100$ , then the system of linear equations has

(Q)  $x = \frac{10}{9}$ ,  $y = -\frac{1}{9}$ ,  $z = 0$  as a solution

(III) If  $\frac{p}{q} \neq 10$ , then the system of linear equations has

(R) infinitely many solutions

(IV) If  $\frac{p}{q} = 10$ , then the system of linear equations has

(S) no solution

(T) at least one solution

The correct option is:

(A) (I)  $\rightarrow$  (T); (II)  $\rightarrow$  (R); (III)  $\rightarrow$  (S); (IV)  $\rightarrow$  (T)

(B) (I)  $\rightarrow$  (Q); (II)  $\rightarrow$  (S); (III)  $\rightarrow$  (S); (IV)  $\rightarrow$  (R)

(C) (I)  $\rightarrow$  (Q); (II)  $\rightarrow$  (R); (III)  $\rightarrow$  (P); (IV)  $\rightarrow$  (R)

(D) (I)  $\rightarrow$  (T); (II)  $\rightarrow$  (S); (III)  $\rightarrow$  (P); (IV)  $\rightarrow$  (T)



$$\left. \begin{aligned} x + y + z &= \underline{1} \\ 10x + 100y + 1000z &= \underline{0} \\ \frac{x}{p} + \frac{y}{q} + \frac{z}{r} &= \underline{0} \end{aligned} \right\}$$

$$D = \begin{vmatrix} 1 & 1 & 1 \\ 10 & 100 & 1000 \\ A+9D & A+99D & A+999D \end{vmatrix} = \begin{vmatrix} 1 & 0 & 0 \\ 10 & 90 & 900 \\ A+9D & 90D & 900D \end{vmatrix} = 0$$

$$D_1 = \begin{vmatrix} 1 & 1 & 1 \\ 0 & 100 & 1000 \\ 0 & A+99D & A+999D \end{vmatrix} = \underline{900}(D-A)$$

$$D_2 = \begin{vmatrix} 1 & 1 & 1 \\ 10 & 0 & 1000 \\ A+9D & 0 & A+999D \end{vmatrix} = 990(D-A)$$



$$D = 0$$

$$D_1 = 900(D - A)$$

$$D_2 = 990(D - A)$$

$$D_3 = 90(D - A)$$

$\infty \text{ Sol}^n$

$$\underline{\underline{D = A}}$$

No Sol<sup>n</sup>

$$\underline{\underline{D \neq A}}$$

③

$$\frac{A + 99D}{A + 9D} \neq 10$$

$$A + 99D \neq 10A + 90D$$

$$9D \neq 9A$$

$$\underline{\underline{D \neq A}}$$

④

$$\underline{\underline{D = A}}$$







## Question Stem

Let  $\alpha, \beta$  and  $\gamma$  be real numbers such that the system of linear equations

$$\begin{aligned}x + 2y + 3z &= \alpha \\4x + 5y + 6z &= \beta \\7x + 8y + 9z &= \gamma - 1\end{aligned}$$

*JEE Adv. 2021*

is consistent. Let  $|M|$  represent the determinant of the matrix

(~~matrix~~ /  $\infty$  Sol<sup>n</sup>)

$$M = \begin{bmatrix} \alpha & 2 & \gamma \\ \beta & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}$$

$$D = 0 \checkmark$$
$$D_1 = D_2 = D_3 = 0$$

3D

Let  $P$  be the plane containing all those  $(\alpha, \beta, \gamma)$  for which the above system of linear equations is consistent, and  $D$  be the square of the distance of the point  $(0, 1, 0)$  from the plane  $P$ .

Q.7 The value of  $|M|$  is 1.

$$|M| = \alpha(1) - \beta(2) - 1(-\gamma)$$
$$= \alpha - 2\beta + \gamma = 1$$

Q.8 The value of  $D$  is 1.5.



$$D_1 = \begin{vmatrix} \alpha & 2 & 3 \\ \beta & 5 & 6 \\ \gamma-1 & 8 & 9 \end{vmatrix} = 0$$

$$\alpha(45-48) - \beta(18-24) + (\gamma-1)(12-15) = 0$$

$$-3\alpha + 6\beta - 3(\gamma-1) = 0$$

$$\alpha - 2\beta + \gamma - 1 = 0$$

$$\underline{\alpha - 2\beta + \gamma = 1}$$

$$(0, 1, 0) \rightarrow \boxed{\alpha - 2\beta + \gamma = 1}$$

$$\left| \frac{0 - 2 + 0 - 1}{\sqrt{1^2 + 2^2 + 1^2}} \right|^2$$

$$\left( \frac{3}{\sqrt{6}} \right)^2$$

$$\frac{9}{6} = \textcircled{1.5}$$





# How to Differentiate a Determinant?



# Differentiation of Determinants



$$F(x) = \begin{vmatrix} f_1(x) & f_2(x) & f_3(x) \\ g_1(x) & g_2(x) & g_3(x) \\ h_1(x) & h_2(x) & h_3(x) \end{vmatrix}$$

$$F'(x) = \begin{vmatrix} f_1'(x) & f_2'(x) & f_3'(x) \\ g_1(x) & g_2(x) & g_3(x) \\ h_1(x) & h_2(x) & h_3(x) \end{vmatrix} + \begin{vmatrix} - & - & - \\ g_1'(x) & g_2'(x) & g_3'(x) \\ - & - & - \end{vmatrix} + \begin{vmatrix} - & - & - \\ - & - & - \\ h_1'(x) & h_2'(x) & h_3'(x) \end{vmatrix}$$



If  $f(x) =$   
 ~~$f'(x)$~~

$$\begin{vmatrix} x+a^2 & ab & ac \\ ab & x+b^2 & bc \\ ac & bc & x+c^2 \end{vmatrix}$$

find  $f'(x)$

$$f'(x) = \begin{vmatrix} 1 & 0 & 0 \\ ab & x+b^2 & bc \\ ac & bc & x+c^2 \end{vmatrix} + \begin{vmatrix} x+a^2 & ab & ac \\ 0 & 1 & 0 \\ ac & bc & x+c^2 \end{vmatrix} + \begin{vmatrix} x+a^2 & ab & ac \\ ab & x+b^2 & bc \\ 0 & 0 & 1 \end{vmatrix}$$





If  $\cancel{a}x^4 + \cancel{b}x^3 + \cancel{c}x^2 + \cancel{d}x + e = \begin{vmatrix} 2x & x-1 & x+1 \\ x+1 & x^2-x & x-1 \\ x-1 & x+1 & 3x \end{vmatrix}$

then the **value of e**, is

A. 0

B. -2

C. 3

D. 2

$x=0$  both side

$$e = \begin{vmatrix} 0 & -1 & 1 \\ 1 & 0 & -1 \\ -1 & 1 & 0 \end{vmatrix}$$



If  $ax^4 + bx^3 + cx^2 + dx + e = \begin{vmatrix} 2x & x-1 & x+1 \\ x+1 & x^2-x & x-1 \\ x-1 & x+1 & 3x \end{vmatrix}$

then the value of d, is

$$4ax^3 + 3bx^2 + 2cx + d = \begin{vmatrix} 2 & 1 & 1 \\ x+1 & x^2-x & x-1 \\ x-1 & x+1 & 3x \end{vmatrix} + \begin{vmatrix} - & - & - \\ 1 & 2x-1 & 1 \\ - & - & - \end{vmatrix} + \begin{vmatrix} - & - & - \\ - & - & - \\ 1 & 1 & 3 \end{vmatrix}$$

$$x=0$$

$$d = \underbrace{\begin{vmatrix} 2 & 1 & 1 \\ 1 & -1 & -1 \\ -1 & 1 & 0 \end{vmatrix} + \begin{vmatrix} - & - & - \\ 1 & -1 & 1 \\ - & - & - \end{vmatrix} + \begin{vmatrix} - & - & - \\ - & - & - \\ 1 & 1 & 3 \end{vmatrix}}$$







# # Cayley – Hamilton Theorem



# Cayley - Hamilton Theorem



x Every square matrix satisfies a specific polynomial equation known as characteristic equation.

$$P(\lambda) = |A - \lambda I|$$

$$P(A) = 0$$

$$|A - \lambda I| = 0$$

$A \rightarrow$  any Sq matrix  $\Rightarrow$  characteristic Eqn

Every Sq matrix will satisfy in its Char. Eqn.



Using **Cayley Hamilton theorem**, find  $A^{-1}$  and  $A^4$

$$A = \begin{bmatrix} 1 & 2 & 0 \\ 2 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$(A - \lambda I) = 0$$

$$\begin{vmatrix} 1-\lambda & 2 & 0 \\ 2 & -1-\lambda & 0 \\ 0 & 0 & 1-\lambda \end{vmatrix} = 0$$

$$(1-\lambda) \{ (-1-\lambda)(1-\lambda) \} - 2(2-2\lambda) = 0$$

$$(1-\lambda)(-1-\lambda+\lambda+\lambda^2) - 4 + 4\lambda = 0$$

$$(1-\lambda)(-1+\lambda^2) - 4 + 4\lambda = 0$$

$$-1 + \lambda^2 + \lambda - \lambda^3 - 4 + 4\lambda = 0$$

$$\lambda^3 - \lambda^2 - 5\lambda + 5 = 0$$

char Eqn

$$CH: \quad \underline{A^3 - A^2 - 5A + 5I = 0}$$

CH  
Kalo?

$$\begin{cases} A^5 & A^{10} & A^{20} & - ? \\ A^{-1} = f(A) \end{cases}$$

$$A^3 = A^2 + 5A - 5I$$

$$A^4 = A A^3$$

$$= A (A^2 + 5A - 5I)$$

$$= A^3 + 5A^2 - 5A$$

$$= A^2 + \cancel{5A} - 5I + 5A^2 - \cancel{5A}$$

$$\boxed{A^4 = 6A^2 - 5I}$$

$$\underline{\underline{A A A A^{-1}}}$$

$$(A^3 - A^2 - SA + SI = 0) A^{-1}$$

$$A^2 - A - SI + S(A^{-1}) = 0$$

$$A^{-1} = \frac{-A^2 + A + SI}{S}$$





Let  $A = \begin{pmatrix} 4 & -2 \\ \alpha & \beta \end{pmatrix}$

$$|A| = 4\beta + 2\alpha = 18$$

If  $A^2 + \gamma A + 18I = O$ , then  $\det(A)$  is equal to

A. -18

☒ B. 18

C. -50

D. 50

$$\begin{vmatrix} 4-\lambda & -2 \\ \alpha & \beta-\lambda \end{vmatrix} = 0$$

$$(4-\lambda)(\beta-\lambda) + 2\alpha = 0$$

$$4\beta - 4\lambda - \lambda\beta + \lambda^2 + 2\alpha = 0$$

$$\lambda^2 + (-4-\beta)\lambda + (2\alpha + 4\beta) = 0$$

CH:  $A^2 + (-4-\beta)A + (2\alpha + 4\beta)I = O$

$$A^2 + \gamma A + 18I = O$$

*JEE Main 2022*

$$-4-\beta = \gamma$$

$$\underline{2\alpha + 4\beta = 18}$$







Let  $A = \begin{pmatrix} 1 & 2 \\ -2 & -5 \end{pmatrix}$ . Let  $\alpha, \beta \in \mathbb{R}$  be such that

$\alpha A^2 + \beta A = 2I$ . Then  $\alpha + \beta$  is equal to -

*JEE Main 2022*

A. -10

B. -6

C. 6

☒ D. 10

$$\begin{vmatrix} 1-\lambda & 2 \\ -2 & -5-\lambda \end{vmatrix} = 0$$

$$(1-\lambda)(-5-\lambda) + 4 = 0$$

$$\lambda^2 + 4\lambda - 1 = 0$$

$$2(A^2 + 4A - I = 0)$$

$$\begin{aligned} 2A^2 + 8A &= 2I \\ \alpha A^2 + \beta A &= 2I \end{aligned}$$

$$\underline{\alpha + \beta = 10}$$



$$(A^2 + 4A = I) A^{-1}$$

$$\underline{A + 4I = A^{-1}}$$



Let  $A = \begin{bmatrix} 1 & 2 \\ -1 & 4 \end{bmatrix}$  If  $A^{-1} = \alpha I + \beta A$ ,  $\alpha, \beta \in \mathbb{R}$ ,  $I$  is a  $2 \times 2$  identity

matrix, then  $4(\alpha - \beta)$  is equal to :

A. 5

B.  $8/3$

C. 2

D. 4

*JEE Main 2021*



# Join with us in Telegram

TELEGRAM CHANNEL "Trigo"


✓ [t.me/unacademyatoms](https://t.me/unacademyatoms)

COMPLETE NOTES AND LECTURES

✓ [tinyurl.com/unacademyatoms](https://tinyurl.com/unacademyatoms)

2 pm

1 Nov DPP Test



**BATCH**

HINDI

Igneous Batch for JEE Advanced & Olympiads 2021

Started on Jul 7

Nishant Vora and 1 more

**BATCH**

HINDI

Evolve Batch Course for Class 12th JEE Main and Advanced 2022

Starts on Apr 7

Anupam Gupta and 2 more

**BATCH**

HINDI

Mega Batch Course for Class 12th JEE Main and Advanced 2022

Starts on Apr 6

Narendra Avasthi and 1 more

**BATCH**

HINDI

Enthuse: Class 12th for JEE Main and Advanced 2022

Starts on Apr 14

Amarnath Anand and 2 more

**BATCH**

HINDI

Final Rapid Revision Batch for JEE Main 2021

Starts on Apr 6

Manoj Chauhan and 2 more

**LIVE**

HINDI PHYSICS

Course of 12th syllabus Physics for JEE Aspirants 2022: Part - I






Lesson 1 • Apr 2, 2021 12:30 PM

D C Pandey


If you want to be the **BEST**  
“Learn” from the **BEST**




# Vector  
# PYQ - Mid Nov  
# CC





## Step 1



## Step 2







Plus  
Live ✓  
Rec ✓  
CC ✓  
TS ✓  
DCSV ✓

**IIT JEE subscription**

ICONIC **PLUS**

- ✓ India's Best Educators
- ✓ Interactive Live Classes
- ✓ Structured Courses & PDFs
- ✓ Live Tests & Quizzes
- ✗ 1:1 Live Mentorship
- ✗ Live Doubt Solving

Duration	Price	EMI	Discount
30 months	₹1,980/mo	No cost EMI	+10% OFF ₹59,399
24 months	₹2,310/mo	No cost EMI	+10% OFF ₹55,440
12 months	₹3,176/mo	No cost EMI	+10% OFF ₹38,115
6 months	₹4,620/mo	No cost EMI	+10% OFF ₹27,720
3 months	₹5,775/mo	No cost EMI	+10% OFF ₹17,325

To be paid as a one-time payment

**NVLIVE**

**IIT JEE subscription**

**PLUS** ICONIC

- ✓ India's Best Educators
- ✓ Interactive Live Classes
- ✓ Structured Courses & PDFs
- ✓ Live Tests & Quizzes
- ✓ **Personal Coach**
- ✓ **Study Planner**

Duration	Price	EMI	Discount
24 months	₹3,750/mo	No cost EMI	+10% OFF ₹90,000
18 months	₹4,000/mo	No cost EMI	+10% OFF ₹72,000
12 months	₹4,875/mo	No cost EMI	+10% OFF ₹58,500
6 months	₹5,700/mo	No cost EMI	+10% OFF ₹34,200

To be paid as a one-time payment

**NVLIVE**

Sm.





# ICONIC PLUS



## Personal Guidance

Get one on one guidance from top exam experts



## Study Planner

Customized study plan with bi-weekly reviews




Live Classes



Weekly Tests



Structured Courses



Unlimited Access



## Test Analysis

Get one on one guidance from top exam experts



## Study Material

Specialised Notes & Practice Sets



## Experts' Guidelines

Study booster workshops by exam experts

# Thank You!

+ SUBSCRIBE



PDF



Download Now !